

Dilaton Field and Massless Particle for 2d Gravity

George Chechelashvili and George Jorjadze

*Dept. of Theoretical Physics, Razmadze Mathematical Institute
Tbilisi, Georgia*

Abstract

It is shown that the model of 2d dilaton gravity is equivalent to the dynamical system of massless particles in the Liouville field.

I Introduction

The Einstein theory of gravity in 2-dimensions is degenerated [1]. The corresponding action

$$S_0 = \int d^2x \sqrt{-g} R \quad (1.1)$$

is expressed only through the ‘surface terms’ and its variation identically vanishes. As a result, the Einstein equations $R_{ab} - 1/2 R g_{ab} = 0$ in 2-dimensions are identities rather than dynamical equations.

The covariant equations $R_{ab} - 1/2 \Lambda g_{ab} = 0$, where Λ is a constant, can be considered as a model of 2d general relativity. These equations are equivalent to $R = \Lambda$, but construction of the corresponding covariant action only in terms of the metric tensor g_{ab} is problematic.

Locally, any 2d Lorentzian manifold is conformally flat [2], i.e. in some local coordinates the metric tensor can be written in the form

$$g_{ab}(X) = \exp q(X) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (1.2)$$

where $X := (x^0, x^1)$ and $q(X)$ is a field. The constant curvature manifold in the conformal gauge (1.2) is described by the Liouville equation [3]

$$(\partial_0^2 - \partial_1^2) q(X) + \Lambda e^{q(X)} = 0, \quad (1.3)$$

which is a variational equation for the functional

$$S[q] = 1/2 \int d^2x \left[(\partial_0 q)^2 - (\partial_1 q)^2 - 2\Lambda e^q \right]. \quad (1.4)$$

Action (1.4) has non-covariant form from the point of view of general relativity and to obtain the corresponding covariant action it is necessary to introduce some additional auxiliary fields. A covariant model with minimal extension is 2d dilaton gravity [4], which arises in Polyakov string theory [5].

For a system with reparametrization invariant action and auxiliary fields, usually, it is difficult to guess what are the physical variables and what kind of physical degrees of freedom they define. Such analysis can be done using the Dirac's procedure for constrained systems based on the Hamiltonian reduction [6]. In the case of 2d dilaton gravity, this procedure eliminates all field degrees of freedom and after full Hamiltonian reduction the system only with finite degrees of freedom remains [7].

In this paper we compare the model of 2d dilaton gravity with the system of massless particle in the corresponding Liouville field [8] and demonstrate their equivalence.

The paper is organized as follows: in Section 2 we prove degeneracy of 2d Einstein gravity. Obtained formulas are used in Section 3 for the description of 2d dilaton gravity. In Section 4 we consider the examples of constant curvature manifolds with global symmetries and analyze the corresponding models of 2d dilaton gravity. In Conclusions we show equivalence of the considered models of 2d gravity and the dynamical systems of massless particles.

II Degeneracy of general relativity in 2-dimensions

The Einstein-Hilbert Lagrangian $\mathcal{L}_0 := \sqrt{-g} R$ can be splitted in two parts [1]

$$\mathcal{L}_0 = \mathcal{L}_1 + \mathcal{L}_2, \quad (2.1)$$

where the first term $\mathcal{L}_1$ contains derivatives of the metric tensor g_{ab} only in the first order

$$\mathcal{L}_1 = \frac{1}{2} \Gamma_{ab}^b \partial_c (\sqrt{-g} g^{ac}) - \frac{1}{2} \Gamma_{ab}^c \partial_c (\sqrt{-g} g^{ab}), \quad (2.2)$$

with

$$\Gamma_{ab}^c := \frac{1}{2} g^{cd} (\partial_a g_{bd} + \partial_b g_{ad} - \partial_d g_{ab}) \quad (2.3)$$

and the second term $\mathcal{L}_2$ is a 'total derivative'

$$\mathcal{L}_2 = \partial_c \left[\sqrt{-g} (g^{ab} \Gamma_{ab}^c - g^{ac} \Gamma_{ab}^b) \right]. \quad (2.4)$$

As a result, $\mathcal{L}_2$ has no contribution in the variational equations $R_{ab} - 1/2 Rg_{ab} = 0$ and the Einstein tensor $G_{ab} := R_{ab} - 1/2 Rg_{ab}$ is defined only through $\mathcal{L}_1$

$$G_{ab} = \frac{1}{\sqrt{-g}} \left(\frac{\partial \mathcal{L}_1}{\partial g^{ab}} - \partial_c \frac{\partial \mathcal{L}_1}{\partial \partial_c g^{ab}} \right). \quad (2.5)$$

Eqs. (2.1)-(2.5) are valid for arbitrary dimension ($D \geq 2$) of spacetime, but the case $D = 2$ posses certain singularity. The Reimann tensor of 2d manifold has only one independent component R_{0101} and the corresponding Einstein tensor G_{ab} identically vanishes for arbitrary metric tensor g_{ab} . Therefore, 2d Einstein equations are identities and not the dynamical equations for g_{ab} . This degeneracy can be seen from the Einstein-Hilbert action (1.1) as well.

Writing the metric tensor of 2d spacetime in the form

$$g_{ab} = \begin{pmatrix} A & B \\ B & C \end{pmatrix} \quad (2.6)$$

and using (2.2)-(2.4) we get

$$\mathcal{L}_1 = -\frac{1}{2G^3} [A(\dot{B}C' - \dot{C}B') + B(\dot{C}A' - \dot{A}C') + C(\dot{A}B' - \dot{B}A')], \quad (2.7)$$

$$\mathcal{L}_2 = \partial_t \left(\frac{\dot{C} - B'}{G} \right) + \partial_x \left(\frac{A' - \dot{B}}{G} \right), \quad (2.8)$$

where

$$G := \sqrt{-g} = \sqrt{B^2 - AC} \quad (2.9)$$

and we also use the notations $x^0 := t$, $x^1 := x$; $\partial_t A := \dot{A}$, $\partial_x A := A'$,

Note that (2.7) can be considered as a value of the 2-form

$$\omega = -\frac{1}{2G^3} (A dB \wedge dC + B dC \wedge dA + C dA \wedge dB), \quad (2.10)$$

on the tangent vectors $V_t := (\dot{A}, \dot{B}, \dot{C})$ and $V_x := (A', B', C')$:

$$\mathcal{L}_1 = \omega(V_t, V_x) \quad (2.11)$$

Since 2-form (2.10) is exact $\omega = d\theta$ [9], (2.11) can be written as

$$\mathcal{L}_1 = \partial_t \theta(V_x) - \partial_x \theta(V_t). \quad (2.12)$$

Thus, in 2-dimensions, $\mathcal{L}_1$ is also a total derivative.

The 1-form θ we can choose by

$$\theta = \frac{1}{G} \left(\frac{B}{C} dC - dB \right) \quad (2.13)$$

and finally, from (2.1), (2.8) and (2.12), we get

$$\sqrt{-g}R(g_{ab}) = \partial_t \left[\frac{1}{G} \left(\dot{C} - 2B' + B \frac{C'}{C} \right) \right] + \partial_x \left[\frac{1}{G} \left(A' - B \frac{\dot{C}}{C} \right) \right]. \quad (2.14)$$

III The model of 2d gravity with dilaton field

Let us consider a model of 2d gravity defined by the action [4]

$$S = \int d^2x \sqrt{-g} \left[\frac{1}{2} g^{ab} \partial_a \phi \partial_b \phi - \phi R + \Lambda \right], \quad (3.1)$$

where ϕ is a scalar (dilaton) field and Λ is a ‘cosmological’ constant.

Variation of (3.1) with respect to the dilaton field defines the dynamical equation

$$-\frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \phi} = \nabla^2 \phi + R = 0. \quad (3.2)$$

To calculate variations of (3.1) with respect to the gravitational field g_{ab} we use the analysis of the Einstein-Hilbert Lagrangian $\mathcal{L}_0 := \sqrt{-g} R(g_{ab})$ given in the previous section. Using (2.1)-(2.4) and neglecting the boundary terms, the functional $S_d := \int d^2x \sqrt{-g} \phi R$ can be written as

$$S_d \simeq \int d^2x (\phi \mathcal{L}_1 - \partial_a \phi X^a), \quad \text{with} \quad X^a = \sqrt{-g} (g^{bc} \Gamma_{bc}^a - g^{ab} \Gamma_{bc}^c). \quad (3.3)$$

The variation of (3.3) reads

$$\frac{1}{\sqrt{-g}} \frac{\delta S_d}{\delta g^{ab}} = \phi G_{ab} + (g_{ab} \nabla^2 - \nabla_a \nabla_b) \phi, \quad (3.4)$$

where $\nabla_a \nabla_b \phi = \partial_{ab}^2 \phi + \Gamma_{ab}^c \partial_c \phi$.

Eq. (3.4) valid for arbitrary dimension of spacetime. Since in 2-dimensions the Einstein tensor G_{ab} vanishes, for the variational equations of (3.1) we obtain

$$\frac{2}{\sqrt{-g}} \frac{\delta S}{\delta g^{ab}} = \partial_a \phi \partial_b \phi + g_{ab} \left(-\frac{1}{2} g^{cd} \partial_c \phi \partial_d \phi - \Lambda \right) - 2(g_{ab} \nabla^2 - \nabla_a \nabla_b) \phi = 0. \quad (3.5)$$

Note that eq.(3.4) in 2-dimensions can be verified from (2.14) as well.

Taking trace of (3.5) and using (3.2) we get that the dynamical system (3.1) describes a spacetime manifold with a constant curvature $R = \Lambda$.

The conformal gauge (1.2) leads to the following form of the dynamical equations (3.2) and (3.5)

$$\partial_{+,-}^2 \varphi = 0, \quad 4\partial_{+,-}^2 q = \Lambda e^q. \quad (3.6)$$

$$(\partial_+ \varphi)^2 - 2\partial_{+,+}^2 \varphi = (\partial_+ q)^2 - 2\partial_{+,+}^2 q, \quad (\partial_- \varphi)^2 - 2\partial_{-,-}^2 \varphi = (\partial_- q)^2 - 2\partial_{-,-}^2 q, \quad (3.7)$$

where $x^\pm := x^1 \pm x^0$ are the light cone coordinates and $\varphi := q - \phi$.

As it was expected, the field $q(X)$ satisfies the Liouville equation (1.3), while $\varphi(X)$ is a ‘free’ field. According to (3.7), the traceless energy-momentum tensors [10-11] of these two fields are equal to each other.

The general solution of the Liouville equation has the form [3]

$$q(x^+, x^-) = \log \frac{8A^{+ \prime}(x^+)A^{- \prime}(x^-)}{|\Lambda| [A^+(x^+) + \epsilon A^-(x^-)]^2} \quad (3.8)$$

where ϵ is a sign of Λ ($\epsilon = \Lambda/|\Lambda|$) and $A^\pm$ are any monotonic functions with $A^\pm \prime > 0$.

The conformal form of the metric tensor (1.2) fixes the gauge freedom only up to the conformal transformations

$$x^\pm \mapsto \tilde{x}^\pm; \quad x^\pm = f^\pm(\tilde{x}^\pm) \quad (3.9)$$

The corresponding transformation of the field q is given by

$$q(x^+, x^-) \mapsto \tilde{q}(x^+, x^-) = q(f^+(x^+), f^-(x^-)) + \log f^{+ \prime}(x^+)f^{- \prime}(x^-). \quad (3.10)$$

The general solution (3.8) can be obtained by the conformal transformation (with $f^\pm(x^\pm) = A^\pm(x^\pm)$) of the Liouville field

$$q(x^+, x^-) = \log \frac{8}{|\Lambda|(x^+ + \epsilon x^-)^2}. \quad (3.11)$$

Hence, all Liouville fields (with fixed Λ) are related to each other by the conformal transformations (3.9)-(3.10). Since the considered system is reparametrization invariant, one can fix the field $q(X)$ by suitable choice of local coordinates.

After fixing the Liouville field $q(X)$ the dynamical freedom of the system is described by the free field $\varphi(X)$, which should be defined from (3.7). These equations are ordinary differential equations for two chiral parts $\varphi_\pm(x^\pm)$ of the free field $\varphi(X) = \varphi_+(x^+) + \varphi_-(x^-)$. For a fixed right hand side of (3.7) the corresponding freedom is described by a finite number of integration constants.

Thus, the system (3.1) has no physical degrees of freedom for field variables and full gage fixing leads to a finite dimensional system. This result can be obtained by the Hamiltonian reduction procedure as well (see [7]).

It should be noted that the conjecture about the conformal form of the metric tensor (1.2) is valid only locally. Therefore, constructions of this section have only a local character. To get the global picture of the system one should specify the global properties of spacetime.

IV The examples of 2d dilaton gravity

The one-sheet hyberboloid

$$-(y_0)^2 + (y_1)^2 + (y_2)^2 = m^{-2}, \quad (4.1)$$

embedded in 3-dimensional Minkowski space is the example of Lorentzian manifold with constant curvature [2]. The parameter m ($m > 0$) defines the scalar curvature $R = -2m^2$. It is obvious that the isometry group of the hyperboloid is $SO_{\uparrow}(2.1)$.

Using the parametrization

$$\begin{aligned} y_0 &= -\frac{\cot m\rho}{m}, & y_1 &= \frac{\cos m\theta}{m \sin m\rho}, & y_2 &= \frac{\sin m\theta}{m \sin m\rho}, \\ \text{where } \rho &\in]0, \pi/m[, & \theta &\in [0, 2\pi/m[\end{aligned} \quad (4.2)$$

we get the conformal form (1.2) of the induced metric tensor

$$g_{ab}(\rho, \theta) = \frac{1}{\sin^2 m\rho} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (4.3)$$

which defines the Liouville field (1.3)

$$q(\rho, \theta) = -\log \sin^2 m\rho, \quad (4.4)$$

for $\Lambda = -2m^2$. The parameters ρ and θ can be identified with the spacetime coordinates x^0 and x^1 , respectively. The corresponding light-cone coordinates we denote by $\theta_{\pm}$ ($\theta_{\pm} := \theta \pm \rho$).

The traceless energy momentum tensor for (4.4) is given by

$$T_{++} := (\partial_+ q)^2 - 2\partial_{+,+}^2 q = -m^2, \quad T_{--} := (\partial_- q)^2 - 2\partial_{-,-}^2 q = -m^2. \quad (4.5)$$

and (3.7) defines the following two equations for the chiral $\varphi_+(x^+)$ and anti-chiral $\varphi_-(x^-)$ parts of the free field φ

$$(\varphi'_+)^2 - 2(\varphi''_+) = -m^2, \quad (\varphi'_-)^2 - 2(\varphi''_-) = -m^2. \quad (4.6)$$

After integration we get

$$\varphi = \varphi_+ + \varphi_- = C - \log \sin^2 \frac{m}{2}(\theta_+ - \alpha_+) - \log \sin^2 \frac{m}{2}(\theta_- - \alpha_-), \quad (4.7)$$

where $\alpha_\pm$ and C ($0 \leq \alpha_\pm < 2\pi/m$) are integration constants. Actually, these three parameters describe all freedom of the system.

The general solution (4.7) has two lines of singularity at $\theta = \alpha_\pm \mp \rho$. According to (4.2), the coordinates of singularity lines on the hyperboloid (4.1) satisfy the relations

$$y_0 + y_1 \cos m\alpha_\pm + y_2 \sin m\alpha_\pm = 0, \quad y_1 \sin m\alpha_\pm - y_2 \cos m\alpha_\pm = \mp \frac{1}{m}, \quad (4.8)$$

which define generatrices of the hyperboloid.

The Cauchy data for the field $\varphi(\rho, \theta)$ at the moment $\rho = \rho_0$ is given by two functions

$$\Phi_{\rho_0}(\theta) := \varphi(\rho_0, \theta) \quad \text{and} \quad \Pi_{\rho_0}(\theta) := \partial_\rho \varphi(\rho_0, \theta). \quad (4.9)$$

These functions are singular at the points $\theta = \alpha_\pm \mp \rho_0$. When these points are different ($\alpha_+ - \rho_0 \neq \alpha_- + \rho_0$), behaviour of the functions (4.9) near the singular points is given by (see (4.7))

$$\Phi_{\rho_0}(\theta) \sim \log \frac{4}{m^2(\theta - \alpha_\pm \pm \rho_0)^2}, \quad \Pi_{\rho_0}(\theta) \sim \mp \frac{2}{\theta - \alpha_\pm \pm \rho_0}. \quad (4.10)$$

If ρ_0 is a ‘moment’ of intersection of singularity lines, the singularity points coincide ($\alpha_+ - \rho_0 = \alpha_- + \rho_0 := \theta_0$). In that case $\Pi_{\rho_0}(\theta)$ is regular and

$$\phi_{\rho_0}(\theta) \sim \log \frac{8}{m^2(\theta - \theta_0)^2}.$$

Thus, the character of singularities of the Cauchy data is preserved in dynamics if the singularity lines does not intersect each other. This case correspond to $\alpha_+ = \alpha_-$.

Another example of spacetime manifold with constant negative curvature can be related to the Liouville field (3.11) for $\Lambda < 0$. This field is singular at $x^0 = 0$. If one excludes the singularity line (see Fig.1), the domains $x^0 > 0$ and $x^0 < 0$ can be considered

as two different patches of the hyperboloid (4.1). The corresponding conformal map to the coordinates (ρ, θ) (see (4.2)) can be written as

$$x^\pm = \frac{1}{m} \tan \frac{m}{2}(\theta \pm \rho). \quad (4.11)$$

The traceless energy momentum tensor for the Liouville field (3.11) identically vanishes and (3.7) defines the following free field

$$\varphi = -2 \log M^2 |(x^+ - \beta_+)(x^- - \beta_-)| \quad (4.12)$$

where $\beta_\pm$ and M are the integration constants. The free fields (4.7) and (4.12) are related by the conformal map (4.11) and the case with non intersecting singularities ($\alpha_+ = \alpha_-$) correspond to $\beta_+ = \beta_-$ (see Fig.1).

Let us consider the stripe (see Fig.2)

$$\mathcal{S} := \{(t, x) \mid t \in \mathbf{R}, x \in]0, \pi/m[\}, \quad (4.13)$$

with the metric tensor

$$g_{\mu\nu}(t, x) = \frac{1}{\sin^2 mx} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (4.14)$$

The coordinates x and t are associated with space (x^1) and time (x^0) coordinates, respectively. From (4.14) we get the positive constant curvature $R = 2m^2$. The isometry group of the manifold (4.13)-(4.14) is the universal covering group of $SL(2, \mathbf{R})$.

The Liouville field $q = -\log \sin^2 mx$ defines the traceless energy-momentum tensor given by (4.5) and for the free field φ we again get (4.7) (substituting $\theta_\pm$ by $x^\pm := x \pm t$). The difference is only in the domain of admissible values of the coordinates $x^\pm$. Since the time coordinate t is not restricted, the singularity lines are given by

$$x \pm t - \alpha_\pm = \frac{2\pi}{m} n_\pm, \quad (4.15)$$

where $n_\pm$ are any integers. In general, the corresponding left and right ‘moving’ lines intersect each other. One can avoid intersection of singularities, assuming again $\alpha_+ = \alpha_-$. In that case the free field $\varphi(t, x)$ is singular only on the zigzag line (see Fig.2) and the class of singularities for the Cauchy data is preserved.

V Conclusion

Analysis of 2d dilaton gravity shows its similarity to the system of massless particle in the corresponding gravitational field. As it is shown in [8], the trajectories of massless particle on the hyperboloid (4.1) are given by the generatrices (4.8), and the ‘zigzag’ line of Fig.2 is the trajectory on the stripe (4.13).

Since the dilaton (scalar) field $\phi = q - \varphi$ has the same singularities as the free field φ , we can associate the dilaton field with the massless scalar particles. In the case of hyperboloid the dilaton field describes two massless particles moving in ‘opposite’ directions (Fig.1), while for the stripe there is only one particle oscillating between the edges of ‘universe’ (Fig.2). By this correspondence we have the kinematical equivalence between the ‘field theory’ model and the particle system. The dynamical equivalence can be achieved using the modified energy-momentum tensor. Adding the term $(g_{ab}\nabla^2 - \nabla_a\nabla_b) \exp(1/2 \phi)$ to the traceless tensor (4.5) (see [12]), we get the localized energy-momentum density, which is concentrated at the singularities of the configurations (4.7).

Acknowledgments

One of the authors (G.J.) is indebted to Paul Sorba for the warm hospitality at Annecy LAAP, where part of this work has been done. G.J. is also grateful to W. Piechocki and G. Weigt for helpful discussions.

This work was supported by the grants from: INTAS (96-0482), RFBR (96-01-00344) and the Georgian Academy of Sciences.

References

- [1] B. A. Dubrovin, S.P. Novikov and A.T. Fomenko, *Modern Geometry*, (Nauka, Moscow, 1979).
- [2] J.A. Wolf, *Space of Constant Curvature*, (Publish or Perish, INC, 1974).
- [3] J. Liouville, *J. Math. Pures Appl.* **18**, 71 (1853).
- [4] N. Seiberg, *Random Surfaces and Quantum Gravity*, edited by O. Alvarez et al. (Plenum Press, New York, 1991) pp. 363-395.

- [5] A. Polyakov, Phys. Lett. B **103**, 207 (1981).
- [6] P.A.M. Dirac, *Lectures on Quantum Mechanics*. Belfer Graduate School of Science, (Yeshive University, New York, 1964).
- [7] G. Jorjadze, *Liouville Field Theory and 2d Gravity*. Proceedings of International Seminar ‘Quarks-98’. Suzdal, 1998 (in press).
- [8] G. Jorjadze and W. Piechocki *Massless Particle in 2d Spacetime with Constant Curvature*. Proceedings of International School and Workshop ISPM. Tbilisi, 1998. Phys. Lett. B (in press).
- [9] G. Jorjadze, L. O’Raifeartaigh and I. Tsutsui. Phys.Lett. B, **336**, n 3-4, 388 (1994).
- [10] G. P. Jorjadze, A. K. Pogrebkov and M. C. Polivanov. Teor. Mat. Fiz., **40**, 221 (1979).
- [11] R. Jackiw, *Progress in Quantum Field Theory*, edited by H. Ezawa and S. Kamefuchi (Elsevier Science Publishers B.V., 1986), pp. 83-98;
- [12] G. P. Jorjadze. Teor. Mat. Fiz. **65**, 323- 333, (1985).




Fig.1 ($\Lambda < 0$)




Fig.2 ($\Lambda > 0$)